



SYSTEM OF NARCOTICS IDENTIFICATION

INSTRUCTIONS FOR USE

TEST H

FOR: Methadone

Test Function

1. This reagent system presumptively identifies Methadone.

How to Use Test H

1. Remove clip and insert into the test pack an amount of powdered suspect material that would fit inside this circle. Reseal with clip and tap gently to assure material falls to the bottom of the pouch.
2. With the printed side of the test pack facing you, break the ampules from left to right. Break by squeezing the center of the ampule with the tips of the thumb and forefinger.
3. Break the left ampule. Agitate gently for one minute. No color is expected at this step.
4. Break right ampule and agitate gently. A bright blue solution indicates the presence of Methadone.

Interpretation of Resulting Colors

1. A blue color will appear after the second ampule is broken if Methadone is present in the suspect material.
2. It is advisable that this test be used in conjunction with others in the Polytesting System.

CAUTION

- A. Since these tests contain strong acids, it is suggested that Test F - Acid Neutralizer be used after testing and before disposal.
- B. Before discarding used test packs, remove clip and add one measure of acid neutralizer from Test F.
- C. Add slowly to prevent bubbling over.
- D. DO NOT reseal the test pouch until effervescence has completely stopped.
- E. Reseal pouch with clip and discard in a tamper-free disposal unit.
- F. No attempt should be made to crush the glass particles after the ampoules are broken.
- G. **Antidote:** In case of contact, immediately flush eyes or skin with water for at least 15 minutes while removing contaminated clothing and shoes. If swallowed, do not give emetics. Contact a physician.

WARNING

These tests are NOT designed for use with liquid samples.



13386 International Parkway
Jacksonville, Florida 32218
904.485.1836 FAX: 904.741.5407
© 2105 Safariland, LLC

A Brand of the Safariland Group